

Acute Type 2 Respiratory Failure From Hypokalemic Respiratory Muscle Paralysis Secondary To Distal (Type1) Renal Tubular Acidosis due To Primary Sjogren's Syndrome

Case Report

Richmond R Gomes^{1*}, Habiba Akhter²

¹ Associate Professor, Medicine, Ad-din Women's Medical College Hospital, Bangladesh.

² Clinical Assistant, Medicine, Ad-din Women's Medical College Hospital, Bangladesh.

Abstract

Tubulointerstitial nephritis (TIN) is the main renal involvement associated with primary Sjogren syndrome (pSS). TIN can manifest as distal renal tubular acidosis (RTA), nephrogenic diabetes insipidus, proximal tubular dysfunction, and others. Distal renal tubular acidosis (RTA) may develop in a large population of patients with Sjogren's syndrome (SS), but most of the subjects are asymptomatic. We report a case of 37 year old woman who presented with acute quadriplegia with acute hypercapnic respiratory failure. She had history of recurrent episodes of limb weakness which was attributable to hypokalemia and initially labelled as hypokalemic periodic paralysis. Later on she was found to have metabolic acidosis rather than alkalosis which pointed towards the diagnosis of renal tubular acidosis (RTA) in the absence of apparent gastrointestinal tract loss. Once the diagnosis of RTA was established, an attempt to search the aetiology revealed that she was having primary Sjogren's syndrome (pSS) though she did not have any symptom at the time of diagnosis. She was found positive for ANA, anti-SSA and anti-SSB. Schirmer test confirmed presence of dry eye. A concomitant existence of autoimmune hypothyroidism was a noteworthy association. Continuous potassium replacement produced rapid and complete recovery from quadriplegia and respiratory failure without requirement for mechanical ventilation. Presentation of this case reminds the importance of vigilance while managing a case of recurrent hypokalemia which might be a rare presenting feature of pSS.

Keywords: Distal Renal Tubular Acidosis; Primary Sjogren's Syndrome; Quadriplegia; Hypercapnic Respiratory Failure.

Introduction

Renal tubular acidosis (RTA) is a group of disorders in which renal excretion of acids is reduced, out of proportion to any reduction in glomerular filtration rate. RTA is characterized by hyperchloremic metabolic acidosis with a normal serum anion gap. There are multiple forms of RTA, depending on which aspect of renal acid handling has been affected (Table 1)[1]. In Type 1 or distal RTA (dRTA), the distal nephron does not lower urine pH normally because the collecting ducts permit excessive back diffusion of H⁺ from lumen to blood. Urinary concentration and potassium conservation also tend to be impaired, so polyuria and polydipsia occur. With the stress of an intercurrent illness, acidosis and hypokalemia can be life-threatening[1]. Acute hypokalemic paralysis is an uncommon cause of acute muscle weakness, and the reported cases of quadriplegia and acute respiratory fail-

ure secondary to hypokalemia are rare [2-5]. We report a case of dRTA who presented with hypokalemic muscular paralysis with acute hypercapnic respiratory failure who did not require ventilation because of fast respiratory function improvement by appropriate treatment with rapid potassium infusion.

Case Report

A 37 years old pleasant, married, muslim, Bangladeshi housewife, known case of hypothyroidism (on 100 mcgm of levothyroxine daily) for 16 years, not known to have any other comorbidities presented to us with the complaints of sudden weakness of both upper and lower limbs for 1 day, shortness of breath for 6 hours and unable to speak for 2 hours. Weakness was sudden in onset and increasing and became so severe that she was unable to move without the help of others and her weakness progressed to the point that she was completely bed ridden. All the proximal

*Corresponding Author:

Dr. Richmond Ronald Gomes,
Associate Professor, Medicine, Ad-din Women's Medical College Hospital, Bangladesh.
Tel: 8801819289499
E-mail: rrichi.dmc.k56@gmail.com

Received: October 27, 2022

Accepted: November 22, 2022

Published: November 24, 2022

Citation: Richmond R Gomes, Habiba Akhter. Acute Type 2 Respiratory Failure From Hypokalemic Respiratory Muscle Paralysis Secondary To Distal (Type1) Renal Tubular Acidosis due To Primary Sjogren's Syndrome. *Int J Resp Dis Care Med.* 2022;5(1):59-63.

Copyright: Richmond Ronald Gomes ©2022. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

and distal muscles and both upper and lower limbs were equally involved but there was no muscle pain. Weakness started in both upper and lower limbs simultaneously. On query her attendant informed that she developed her symptoms after taking heavy meal. There is no history of trauma, pain in the neck or dysphagia, sensory complaints like paresthesia. She also developed breathlessness for 6 hours which was sudden and progressively increasing but not associated with chest pain, sputum or haemoptysis. For last two hours she was unable to speak but can understand speech and can follow commands like closing her eyes. She denied any history of fever, headache, visual complaints in the form of blurring of vision, diplopia, convulsion, urinary and fecal incontinence, recent vaccination. There was also no history of recent previous infection like respiratory tract infection, urinary tract infection or diarrhea. Patient's attendant also complained of occasional rash about which they could not elaborate. She has history of similar type of attack about 1 year back which was managed conservatively in a local hospital. There was no similar type of illness in her family. She has three children and all are in good health. She had no history of abortion, no bad obstetric history and all of her pregnancy was uneventful.

On physical examination, she was conscious, oriented, but a bit drowsy with GCS 14/15 and had are flexicquadruplegia. Power was graded 0/5 in the legs and in the arms. Plantar responses were absent bilaterally. All modalities of sensation are intact, sings of meningeal irritation were absent. Cerebellar signs, stance and gait could not be evaluated. She had palpable rash over both lower

limbs (figure 1), buttocks and over both palm. Her pulse was 60 beats/min, blood pressure 110/70 mmHg, respiratory rate 30 breath/min. Her chest was clear on auscultation, oxygen saturation was 97% on 5 L of oxygen. The rest of physical exam was unremarkable.

On initial laboratory investigations, complete blood count, random blood sugar, SGPT, serum creatinine, CPK, chest x ray revealed normal results except ESR was 68 mm in 1st hour. Serum Electrolyte showed sodium 138 mmol/L, potassium 1.7 mmol/L, chloride 111 mmol/L, bicarbonate 16.5 mmol/L. Anion gap was normal (12.2 mmol/L). Urinary potassium was elevated with 101 mmol/L. TSH, serum calcium, serum magnesium level were normal. Electrocardiogram showed sinus bradycardia, diffuse ST depression, T inversion, U wave in precordial limbs and prolonged PR interval. (Figure 2)

Arterial blood gas (ABG) analysis revealed mixed respiratory and metabolic acidosis with hypercapnic respiratory failure (pH 7.03, PO₂-69 mm, PCO₂- 64 mm, HCO₃-16 mmol/L). She was started treatment with intravenous sodium bicarbonate along with intravenous and oral potassium chloride and showed dramatic clinical improvement in 1 day. On next day, her muscle power improved to 4/5 in both upper and lower limbs, respiratory rate was 16 breaths/min and can speak well. SaO₂ was 96% in room air. Repeat ABG in room air revealed resolved hypercapnic respiratory failure (pH 7.40, PO₂-88 mm, PCO₂- 30 mm, HCO₃-14 mmol/L). Repeat serum electrolyte revealed sodium

Figure 1. Showing Palpable Vasculitic Rash Over Lower Limb.




Figure 2. Electrocardiogram showing sinus bradycardia, diffuse ST depression, T inversion (black arrow), U wave (red arrow) in precordial limbs and prolonged PR interval.



Figure 3. Positive Schirmer test showing 1 mm wet on filter paper after 5 minutes.



147 mmol/L, potassium 2.9 mmol/L, chloride 109 mmol/L and bicarbonate 18 mmol/L. Despite of correction of respiratory acidosis her metabolic acidosis persisted. Urine routine examination showed pH 7.2, trace proteinuria without any RBC, pus cell or any casts. So type 1 or distal renal tubular acidosis (dRTA) was suspected and for further evaluation of underlying cause for dRTA, on query, she revealed that she has been suffering from occasional rash for last 5 years which was more prominent in her legs and buttock area and also involved her palm sparing her trunk and face. Rashes care initially itchy and then became painful. She developed these rashes occasionally for last 5 years and resolved spontaneously. She denied any history of joint pain, oral ulcer, muscle pain, hair fall, and weight loss. She also complained of dryness of mouth for last 15 years, that was increasing and became so severe that she cannot swallow her food without water. She also developed dry eyes for same duration and there is no tear during her emotional event like crying. Bedside Schirmer's test was done and found positive(1 mm wet on filter paper in 5 minutes- figure 3). She also complaints of bluish discoloration of her fingers and pain on cold exposure and she has been suffering from fatigue and weakness for last 15 years and dry cough and worsening shortness of breath on exertion for last 2 years. With these symptoms, she visited a number of physicians including ophthalmologists and was managed symptomatically without any improvement.

Further investigations revealed: ANA was strongly positive on Hep2 cell with fine speckled pattern. Anti dS DNA, lupus anti coagulant, anti cardiolipin antibody was negative. ENA profile showed positive anti SSA(Ro) and anti SSB(La). RA test was positive in high titer, Serum total protein was 77 gm/L, albumin 35 gm/L and globulin 42 gm/L, albumin globulin ratio was 1: 1.42. Anti TPO was > 1300 U/ml(positive >60 U/ml). C4 level was reduced to 0.1 g/L(normal 0.15-0.45 g/L). Protein Electrophoresis showed polyclonal hypergammaglobulinemia with prominent IgG and IgM. Pulmonary function tests showed mild restrictive defect as evidenced by low FEV1(2.02 L) and low FVC(2.05L) with normal FEV1/FVC (94.5%). DLCO was moderately reduced with 55% of predicted value. KCO was normal suggestive of parenchymal restrictive disease. Patient refused for lip biopsy. So final diagnosis of recurrent hypokalemic paralysis due to distal (type 1) renal tubular acidosis secondary to primary Sjogren's syndrome with type 3 cryoglobulinemic vasculitis with autoimmune hypothyroidism was made. She was discharged with sodium

bicarbonate, potassium citrate, hydroxychloroquine (400 mg daily) and artificial tear. After 1 month, on out patient door follow up she was doing well.

Discussion

Approximately, 98% of total body potassium is located intracellularly. Sixty percent of intracellular potassium is within skeletal muscle, which may explain the predominance of muscular symptoms in disorders producing hypokalemia [6, 7]. Although, the effects of hypokalemia are multisystemic, potassium deficiency most seriously affects the neuromuscular system, and symptoms related to hypokalemia are typically muscular. Serum potassium concentrations of 3.0 to 3.5 mmo U-1 may be associated with mild muscle weakness and myalgia. Serum potassium concentrations of 2.5 to 3.0 mmol/L are associated with the development of clinically significant muscle weakness. The muscle weakness generally is limited to the limbs and limb girdles. The respiratory or cranial musculature are rarely involved and characteristically spared in muscle weakness secondary to hypokalemia. When severe, hypokalemia can impair respiratory function leading to hypoventilation, and acute respiratory failure [2-5, 8, 9]. Hypokalemic muscleweakness may simulateGuillain -Barre syndrome [10-12]. When the serum potassium level fallsbelow 2.5 mmol/L, rhabdomyolysis may occur [7, 13, 14]. The lowest serum potassium level reported was 1.1 mmol./L this was associated with areflexic quadriplegia, coma, and acute respiratory failure managed with IPPV8.

Hypokalemia is a common clinical problem, and is the most frequent electrolyte disorder encountered in clinical practice. The major causes of hypokalemia are: Inadequate potassium intake, excessive skin, renal (diuretics, adrenal insufficiency, salt losing nephropathy), orgastrointestinal losses (diarrhoea, vomiting), and increased entry into cells by a variety of mechanisms including increased availability of insulin, elevated beta adrenergic activity and elevation in extracellular pH (metabolic orrespiratory alkalosis).

RTA is characterized by hyperchloremic metabolic acidosis with a normal serum anion gap. Thereare multiple forms of RTA, depending on which aspect of renal acid handling has been affected (Table 1).

The presence of severe hypokalemia (K+ 1.7mmol/L) with nor-

Table 1. Comparison of different forms of renal tubular acidosis.

Finding	Type 1 RTA	Type 2 RTA	Type 4 RTA
Normal anion gap acidosis	Yes	Yes	Yes
Minimum urine pH	>5.5	<5.5	<5.5
% filtered bicarbonate excreted	<10	>15	<10
Serum potassium	Low	Low	High
Stones/ nephrocalcinosis	Yes	No	No
Daily acid excretion	Low	Normal	Low
Urine anion gap	Positive	Positive	Positive
Daily bicarbonate replacement needs	<4 mmol/kg	>4 mmol/kg	<4 mmol/kg

mal anion gap metabolic acidosis and urinary pH >5.5 suggested the diagnosis of Type 1 dRTA. Although quadriplegia is a well-known complication of severe hypokalemia, acute respiratory failure due to dRTA is rarely reported [15]. A thorough search of the available literature (Medline) revealed only six cases of dRTA leading to hypokalemic respiratory failure requiring mechanical ventilation [16-21].

Ascending paralysis of the extremities suggested Guillain-Barré syndrome, however the presence of hypokalemia, metabolic acidosis and an alkaline urine confirmed the diagnosis, and lumbar puncture was not performed. Moreover, there was a significant improvement in muscle power after K⁺ supplementation.

The pathogenesis of classic hypokalemic RTA is not yet clear. The hallmark is the inability to acidify the urine to pH <5.5; there is no impairment in bicarbonate reabsorption in proximal tubules [1]. In the majority of patients, there is a defect in H⁺ secretion by the H⁺-K⁺-ATPase pump in tubular cells which causes hypokalemia, decrease in NH₄⁺ excretion, hyperchloaemic metabolic acidosis and volume depletion. Occasionally, strikingly severe hypokalemia, metabolic acidosis and hypocalcaemia may require immediate therapy. This constellation of findings in an emergency setting has been labelled as a “crisis of dRTA” [22]. Clinical presentation in our patient was also quite catastrophic, with quadriplegia, acute respiratory failure which could have proved fatal without prompt and effective management.

Chronic positive acid balance causes Ca²⁺, Mg²⁺ and PO₄³⁻ wasting and there is an increased incidence of nephrocalcinosis. A prominent feature of dRTA is abnormal Ca²⁺ metabolism leading to musculoskeletal complaints. Hypercalciuria, alkaline urine and low levels of urinary citrate result in calcium phosphate stones [22].

The majority of patients with dRTA have it in association with a systemic illness such as Sjögren's syndrome, hypergammaglobulinaemia, chronic active hepatitis (CAH) or lupus. The frequency of dRTA in Sjögren's syndrome has been reported to be 25 to 40% [22]. Ohtani et al and Poux et al have described hypokalemic quadriplegia and respiratory arrest due to dRTA in patients with primary Sjögren's syndrome [18, 19]. Koul and Saleem reported a case of CAH with dRTA which presented with hypokalemic muscular paralysis requiring respiratory assistance [21].

The diagnostic criteria of Sjogren's syndrome were satisfied by the findings of xerostomia, high titer of anti SSA and anti SSB antibodies. The frequency of dRTA in Sjogren's syndrome has been reported to be about 25-40%. [23, 24]. There are many case reports of Sjogren's syndrome associated with hypokalemic quadriplegia [25-31]. Tsuboi et al described that the periodic paralysis was observed in almost 40% of Sjogrens syndrome cases associated with dRTA [31]. However only three cases were reported that respiratory arrest depends on severe hypokalemia associated with renal tubular acidosis due to various causes [32-34] and there is only one case report describing respiratory failure due to Sjogren's syndrome [34]. There is no significant difference at the level of serum potassium between respiratory arrest group and quadriplegia alone group (1.0-2.8 mmol/L). These results suggests that the progress for respiratory arrest may be may be influenced by interindividual differences in the sensitivity towards hypokalemia (and presumably also transmembrane K gradient) of respiratory

muscles. Even patients with hypokalemia <1.7 mEq/l were liable to develop respiratory arrest. The above cases, including our own patient, were all female, and showed good prognosis with immediate respiratory support and potassium supplementation. However, Nimmannit et al. described fatal cases of hypokalemic respiratory failure and ventricular fibrillation due to endemic RTA in Thailand. This condition occurred in otherwise healthy young males and sometimes resulted in nocturnal death which is different from the present case.

Treatment of dRTA involves correction of metabolic acidosis by administration of alkali in an amount sufficient to neutralize the production of diet-derived acids. This is usually equal to 1-3 mmol/kg/.day of sodium bicarbonate in adults. In children, a large amount of bicarbonate must be administered to correct acidosis and maintain normal growth. Sustained correction of metabolic acidosis usually suppresses renal Na⁺ and K⁺ wasting with attendant correction of extracellular fluid volume and hypokalemia. Therefore in most patients with dRTA, K⁺ supplementation is not necessary. However, if the patient presents with hypokalemic respiratory paralysis, the K⁺ deficit should be replaced without delay before initiating bicarbonate therapy [22], because if bicarbonate is given alone it may further worsen the already existing hypokalemia. The rate of infusion of KCl should be up to a maximum of 0.25 mmol/kg/.hr. These patients are often volume depleted and treatment should include correction of the fluid deficit if necessary guided by central venous pressure monitoring.

It is well known that hypokalemia can cause muscle weakness and respiratory failure, but is probably under-appreciated. A prompt diagnosis and treatment of hypokalemia is warranted to prevent respiratory failure and the requirement for mechanical ventilation, and to prevent other complications, mainly cardiac.

Conclusion

This case report highlights the fact that the dRTA with severe hypokalemia should be kept in mind in any patient presenting with respiratory failure. Secondly, all patients with dRTA must be investigated for any associated systemic illnesses and put on life-long alkalinizing agents. In pSS, RTA is commonly asymptomatic. However, if such a state persists, patient may develop quadriplegia. Furthermore, if adequate treatment is not received, muscle paralysis may progress to respiratory arrest. Although respiratory arrest associated with Sjogren's syndrome is very rare, this complication is very severe and can be fatal. It is important to pay attention to the occurrence of severe hypokalemia with metabolic acidosis and provide adequate treatment for this combination in patient with pSS.

References

- [1]. Asplin JR, Coe FR. Hereditary Tubular Disorders. In: Braunwald E, Fauci AS, Kasper DL et al, eds. Harrison's Principles of Internal Medicine. New York: McGraw-Hill; 2001.
- [2]. Mutlu GM, Factor P. Acute-onset quadriplegia, respiratory failure, and ventricular tachycardia in a 21-year-old man following a soccer match. *Chest*. 2002 Jun;121(6):2036-9. PubMed PMID: 12065374.
- [3]. Davies RG, Gemmell L. Severe hypokalaemia causing acute respiratory failure. *Anaesthesia*. 2001 Jul;56(7):694-5. PubMed PMID: 11437788.
- [4]. Chandrashekar R, Claxton AR. Acute respiratory failure secondary to severe hypokalaemia. *Anaesthesia*. 2000 Dec;55(12):1221-2. PubMed PMID:

- 11121946.
- [5]. Dunn SR, Farnsworth TA, Karunaratne WU. Hypokalaemic, hyperchloraemic metabolic acidosis requiring ventilation. *Anaesthesia*. 1999 Jun;54(6):566-8. PubMed PMID: 10403871.
 - [6]. Knochel JP. Neuromuscular manifestations of electrolyte disorders. *Am J Med*. 1982 Mar;72(3):521-35. PubMed PMID: 7036740.
 - [7]. Riggs JE. Neurologic manifestations of electrolyte disturbances. *NeurolClin*. 2002 Feb;20(1):227-39, vii. PubMed PMID: 11754308.
 - [8]. Latière M, Dumont JC, Olmer M, François G. Hypokalemic quadriplegia and coma in renal tubular acidosis. *Ann Fr Anesth Reanim*. 1989;8(2):133-6. PubMed PMID: 2735571.
 - [9]. Fogliani J, Giraud E, Henriquet D, Maitresse B. Voluntary barium poisoning. *Ann Fr Anesth Reanim*. 1993;12(5):508-11. PubMed PMID: 8311359.
 - [10]. Koley TK, Goyal AK, Gupta MD. Barium poisoning mimicking Guillain-Barré syndrome. *J Assoc Physicians India*. 2001 Jun;49:656-7. PubMed PMID: 11584944.
 - [11]. Warner TT, Mossman S, Murray NM. Hypokalaemia mimicking Guillain-Barré syndrome. *J Neurol Neurosurg Psychiatry*. 1993 Oct;56(10):1134-5. PubMed PMID: 8410018.
 - [12]. Royston A, Prout BJ. Carbenoxolone-induced hypokalaemia simulating Guillain-Barré syndrome. *Br Med J*. 1976 Jul 17;2(6028):150-1. PubMed PMID: 1276838.
 - [13]. Champion DS, Arias JM, Carter NW. Rhabdomyolysis and myoglobinuria. Association with hypokalaemia of renal tubular acidosis. *JAMA*. 1972 May 15;220(7):967-9. PubMed PMID: 5067452.
 - [14]. Kao KC, Tsai YH, Lin MC, Huang CC, Tsao CY, Chen YC. Hypokalemic muscular paralysis causing acute respiratory failure due to rhabdomyolysis with renal tubular acidosis in a chronic glue sniffer. *J Toxicol Clin Toxicol*. 2000;38(6):679-81. PubMed PMID: 11185978.
 - [15]. Gk A, Mittal VK. Periodic paralysis with renal tubular acidosis. *Neurol India*. 1981;30:188-91.
 - [16]. Tyagi A, Rautela RS, Bhattacharya A. Renal tubular acidosis presenting with respiratory arrest. *Indian J Pediatr*. 2002 Feb;69(2):197. PubMed PMID: 11929040.
 - [17]. Fujimoto T, Shiiki H, Takahi Y, Dohi K. Primary Sjögren's syndrome presenting as hypokalaemic periodic paralysis and respiratory arrest. *Clin Rheumatol*. 2001;20(5):365-8. PubMed PMID: 11642520.
 - [18]. Ohtani H, Imai H, Kodama T, Hamai K, Komatsuda A, Wakui H, et al. Severe hypokalaemia and respiratory arrest due to renal tubular acidosis in a patient with Sjögren syndrome. *Nephrol Dial Transplant*. 1999 Sep;14(9):2201-3. PubMed PMID: 10489232.
 - [19]. Poux JM, Peyronnet P, Le Meur Y, Favereau JP, Charnes JP, Leroux-Robert C. Hypokalemic quadriplegia and respiratory arrest revealing primary Sjögren's syndrome. *Clin Nephrol*. 1992 Apr;37(4):189-91. PubMed PMID: 1582057.
 - [20]. Alvarez J, Low RB. Acute respiratory arrest due to hypokalemia. *Ann Emerg Med*. 1988 Mar;17(3):288-9. PubMed PMID: 3345025.
 - [21]. Koul PA, Saleem SM. Chronic active hepatitis with renal tubular acidosis presenting as hypokalemic periodic paralysis with respiratory failure. *Acta Paediatr*. 1992 Jun-Jul;81(6-7):568-9. PubMed PMID: 1392378.
 - [22]. Du Bose TD Jr, Cogan MG, Rector FC Jr. Acid Base Disorders. In: Brenner BM, ed. *Brenner & Rector's The Kidney*; 5th Ed. Philadelphia: WB Saunders Company; 1996; 929-998.
 - [23]. Shearn MA, Tu WH. Latent renal tubular acidosis in Sjögren's syndrome. *Ann Rheum Dis*. 1968 Jan;27(1):27-32. PubMed PMID: 4170879.
 - [24]. Siamopoulos KC, Mavridis AK, Elisaf M, Drosos AA, Moutsopoulos HM. Kidney involvement in primary Sjögren's syndrome. *Scand J Rheumatol Suppl*. 1986;61:156-60. PubMed PMID: 3473626.
 - [25]. Christensen KS. Hypokalemic paralysis in Sjögren's syndrome secondary to renal tubular acidosis. *Scand J Rheumatol*. 1985;14(1):58-60. PubMed PMID: 4001878.
 - [26]. Nagashima K, Wakayama M, Yaguchi M, Yoshida T, Okamoto K, Hirai S. A patient with Sjögren syndrome with central pontine myelinolysis and hypokalemic myopathy. *Rinsho Shinkeigaku*. 1996 Nov;36(11):1240-4. PubMed PMID: 9046856.
 - [27]. Pun KK, Wong CK, Tsui EY, Tam SC, Kung AW, Wang CC. Hypokalemic periodic paralysis due to the Sjögren syndrome in Chinese patients. *Ann Intern Med*. 1989 Mar 1;110(5):405-6. PubMed PMID: 2916810.
 - [28]. Chang YC, Huang CC, Chiou YY, Yu CY. Renal tubular acidosis complicated with hypokalemic periodic paralysis. *Pediatr Neurol*. 1995 Jul;13(1):52-4. PubMed PMID: 7575850.
 - [29]. Raskin RJ, Tesar JT, Lawless OJ. Hypokalemic periodic paralysis in Sjögren's syndrome. *Arch Intern Med*. 1981 Nov;141(12):1671-3. PubMed PMID: 7305577.
 - [30]. Hattori N, Hino M, Ishihara T, Moridera K, Ikekubo K, Kurahachi H. Hypokalemic paralysis associated with distal renal tubular acidosis. *Intern Med*. 1992 May;31(5):662-5. PubMed PMID: 1504432.
 - [31]. Tsuboi Y, Nijima K, Jyoshita Y, Yoshida M, Muto S. A case of renal tubular acidosis type I RTA with Sjögren's syndrome, hypokalemic periodic paralysis, and selective hypoaldosteronism. *Saishin Gaku*. 1990;45:610-17.
 - [32]. Bridi GS, Falcon PW, Brackett NC Jr, Still WJ, Sporn IN. Glomerulonephritis and renal tubular acidosis in a case of chronic active hepatitis with hyperimmunoglobulinemia. *Am J Med*. 1972 Feb;52(2):267-78. PubMed PMID: 5058508.
 - [33]. Koul PA, Saleem SM. Chronic active hepatitis with renal tubular acidosis presenting as hypokalemic periodic paralysis with respiratory failure. *Acta Paediatr*. 1992 Jun-Jul;81(6-7):568-9. PubMed PMID: 1392378.
 - [34]. Poux JM, Peyronnet P, Le Meur Y, Favereau JP, Charnes JP, Leroux-Robert C. Hypokalemic quadriplegia and respiratory arrest revealing primary Sjögren's syndrome. *Clin Nephrol*. 1992 Apr;37(4):189-91. PubMed PMID: 1582057.
 - [35]. Nimmannit S, Malasit P, Chaovakul V, Susaengrat W, Vasuvattakul S, Nilwarrangkur S. Pathogenesis of sudden unexplained nocturnal death (lai tai) and endemic distal renal tubular acidosis. *Lancet*. 1991 Oct 12;338(8772):930-2. PubMed PMID: 1681278.